

In this lecture, we study: let  $p$  be an odd prime,

$$\left(\frac{2}{p}\right) = ?$$

By Euler's criterion, we have:

$$2^{\frac{p-1}{2}} \equiv \left(\frac{2}{p}\right) \pmod{p}$$

Therefore, we just need to investigate  $2^{\frac{p-1}{2}} \pmod{p}$ .

We look at two examples and then try to generalize it.

Example I:  $p = 13$ . Set  $P = \frac{p-1}{2} = 6$ .

We look at all even integers between 0 and 13.

$$2 \quad 4 \quad 6 \quad 8 \quad 10 \quad 12. \quad (6 \text{ numbers})$$

$$\begin{aligned} \bullet 2 \cdot 4 \cdot 6 \cdot 8 \cdot 10 \cdot 12 &= 2(1) \cdot 2(2) \cdot 2(3) \cdot 2(4) \cdot 2(5) \cdot 2(6) \\ &= 2^6 \cdot (1)(2)(3)(4)(5)(6) \\ &= 2^6 \cdot 6! \end{aligned}$$

$$\begin{aligned} \bullet 2 \cdot 4 \cdot \underset{\Delta}{6} \cdot 8 \cdot 10 \cdot 12 &\equiv 2 \cdot 4 \cdot 6 \pmod{13} \\ &\equiv 2 \cdot 4 \cdot 6 \cdot (-5)(-3)(-1) \pmod{13} \\ &\equiv (-1)^3 \cdot 6! \end{aligned}$$

$$\Rightarrow 2^6 \cdot 6! \equiv (-1)^3 \cdot 6! \pmod{13}$$

$$\Rightarrow 2^6 \equiv -1 \pmod{13} \Rightarrow \left(\frac{2}{13}\right) = -1$$

Example II.  $P=17$   $P = \frac{17-1}{2} = 8$

We look at all even integers between 0 and 17.

2 4 6 8 10 12 14 16 (8 numbers)

$$\bullet 2 \cdot 4 \cdot 6 \cdot 8 \cdot 10 \cdot 12 \cdot 14 \cdot 16 = 2(1) \cdot 2(2) \cdot 2(3) \cdot 2(4) \cdot 2(5) \cdot 2(6) \cdot 2(7) \cdot 2(8)$$
$$= 2^8 \cdot 8! \pmod{17}$$

$$\bullet 2 \cdot 4 \cdot 6 \cdot 8 \cdot 10 \cdot 12 \cdot 14 \cdot 16 \equiv 2 \cdot 4 \cdot 6 \cdot 8 \cdot (10-17)(12-17)(14-17)(16-17)$$
$$\quad \triangle \quad \equiv 2 \cdot 4 \cdot 6 \cdot 8 \cdot (-7)(-5)(-3)(-1) \pmod{17}$$
$$\equiv (-1)^4 \cdot 8! \pmod{17}$$

$$\Rightarrow 2^8 \cdot 8! \equiv (-1)^4 \cdot 8! \pmod{17}$$

$$\Rightarrow 2^8 \equiv 1 \pmod{17} \quad \left(\frac{2}{17}\right) = 1 .$$

We consider a general  $p$ , set  $P = \frac{p-1}{2}$ .

$$\begin{aligned} \cdot 2 \cdot 4 \cdot 6 \cdots (p-1) &= 2(1) \cdot 2(2) \cdot 2(3) \cdots 2 \cdot \left(\frac{p-1}{2}\right) \\ &= 2^{\frac{p-1}{2}} \cdot \left(\frac{p-1}{2}\right)! \end{aligned}$$

$$\cdot 2 \cdot 4 \cdot 6 \cdots (p-1) = 2 \cdot 4 \cdot 6 \cdots \frac{p-1}{2} \cdot \frac{p+3}{2} \cdot \frac{p+7}{2} \cdots (p-1)$$

Here we only considered the case when  $P$  is even.

We can do this in a similar way when  $P$  is odd.

$$\begin{aligned} &\equiv 2 \cdot 4 \cdot 6 \cdots \frac{p-1}{2} \cdot \left(\frac{p+3}{2} - p\right) \left(\frac{p+7}{2} - p\right) \cdots (p+1-p) \pmod{p} \\ &\equiv 2 \cdot 4 \cdot 6 \cdots \frac{p-1}{2} \cdot \left(-\frac{p-3}{2}\right) \left(-\frac{p-7}{2}\right) \cdots (-1) \pmod{p} \\ &\equiv (-1)^* \cdot \left(\frac{p-1}{2}\right)! \pmod{p} \end{aligned}$$

$(*)$  = number of integers in the list

2, 4, 6, 8, ...  $p-1$

that are larger than  $\frac{p-1}{2}$ .

$$\Rightarrow 2^{\frac{p-1}{2}} \cdot \left(\frac{p-1}{2}\right)! \equiv (-1)^* \cdot \left(\frac{p-1}{2}\right)! \pmod{p}$$

$$\Rightarrow 2^{\frac{p-1}{2}} \equiv (-1)^* \pmod{p} \Rightarrow \left(\frac{2}{p}\right) = (-1)^*$$

## Theorem 21.4 (Quadratic Reciprocity, Part II)

Let  $p$  be an odd prime.

$$\left(\frac{2}{p}\right) = \begin{cases} 1 & \text{if } p \equiv 1, 7 \pmod{8} \\ -1 & \text{if } p \equiv 3, 5 \pmod{8} \end{cases}$$

Proof: We need to consider 4 cases. Here we only do

$$p \equiv 1 \pmod{8} \quad \text{and} \quad p \equiv 5 \pmod{8}$$

$$\text{Case: } p \equiv 1 \pmod{8} \quad p = 8k + 1. \quad P = \frac{p-1}{2} = 4k.$$

$$2, 4, 6, \dots, 4k, 4k+2, \dots, 8k \quad (\text{4k numbers})$$

We need to count how many numbers are  $> P = 4k$

That is,  $4k+2, 4k+4, \dots, 8k$ .

There are  $2k$  numbers in total and hence

$$\begin{aligned} 2^{P-1} &= 2^{4k} \equiv (-1)^{2k} \pmod{p} \\ &\equiv 1 \pmod{p} \end{aligned}$$

$$\Rightarrow \left(\frac{2}{p}\right) = 1$$

$$\text{Case: } p \equiv 5 \pmod{8} \quad p = 8k+5 \quad P = \frac{P-1}{2} = 4k+2.$$

We look at

$$2, 4, 6, \dots, 4k+2, \underbrace{4k+4, 4k+6, \dots, 8k+4}_{2k+1 \text{ numbers.}}$$

$$\begin{aligned} \text{Therefore, } 2^{P-1} &= 2^P \equiv (-1)^{2k+1} \pmod{p} \\ &\equiv -1 \pmod{p} \\ \Rightarrow \left(\frac{2}{p}\right) &= -1 \end{aligned}$$

□

**Remark:** In the proof of the theorem, the key observation is:

$$2^{\frac{P-1}{2}} \equiv (-1)^{\#} \pmod{p}$$

where  $\#$  = number of integers in the list

$$2, 4, 6, 8, \dots, P-1$$

that are larger than  $\frac{P-1}{2}$ .

Here is another way to interpret  $2, 4, 6, 8, \dots, P-1$ :

$$2 \cdot 1, 2 \cdot 2, 2 \cdot 3, \dots, 2 \cdot \frac{P-1}{2}.$$

Later, when we consider  $(\frac{a}{p})$ , we will look at:  
 $a \cdot 1, a \cdot 2, a \cdot 3, \dots, a \cdot \frac{P-1}{2}$ .