

Recall: let  $p$  be an odd prime. Then

$$\left(\frac{-1}{p}\right) = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{4} \\ -1 & \text{if } p \equiv 3 \pmod{4} \end{cases}$$

$$\left(\frac{2}{p}\right) = \begin{cases} 1 & \text{if } p \equiv 1, 7 \pmod{8} \\ -1 & \text{if } p \equiv 3, 5 \pmod{8} \end{cases}$$

Theorem 22.1. (Quadratic Reciprocity) Let  $p, q$  be distinct odd primes, then

$$\left(\frac{q}{p}\right) \cdot \left(\frac{p}{q}\right) = (-1)^{\frac{p-1}{2} \cdot \frac{q-1}{2}}$$

Remark: (1) If  $p \equiv 1 \pmod{4}$  or  $q \equiv 1 \pmod{4}$ , then

$$\frac{p-1}{2} \cdot \frac{q-1}{2} \text{ is even.}$$

$$\text{Therefore, } \left(\frac{q}{p}\right) \cdot \left(\frac{p}{q}\right) = 1 \Rightarrow \left(\frac{q}{p}\right) = \left(\frac{p}{q}\right)$$

(2) If  $p \equiv 3 \pmod{4}$  and  $q \equiv 3 \pmod{4}$ , then

$$\frac{p-1}{2} \cdot \frac{q-1}{2} \text{ is odd}$$

$$\text{Therefore, } \left(\frac{q}{p}\right) \cdot \left(\frac{p}{q}\right) = -1 \Rightarrow \left(\frac{q}{p}\right) = -\left(\frac{p}{q}\right)$$

Recall: Let  $p$  be an odd prime, and  $a, b$  be integers.

(1) Multiplication Rule:

$$\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right)\left(\frac{b}{p}\right)$$

(2) Congruence Rule: if  $a \equiv b \pmod{p}$

$$\left(\frac{a}{p}\right) = \left(\frac{b}{p}\right)$$

Example: Find  $\left(\frac{-14}{137}\right)$

Solution:  $14 = (-1) \times 2 \times 7 \Rightarrow \left(\frac{-14}{137}\right) = \left(\frac{-1}{137}\right)\left(\frac{2}{137}\right)\left(\frac{7}{137}\right)$

$$(1) \quad 137 \equiv 1 \pmod{4} \Rightarrow \left(\frac{-1}{137}\right) = 1$$

$$(2) \quad 137 \equiv 1 \pmod{8} \Rightarrow \left(\frac{2}{137}\right) = 1$$

$$(3) \quad 7 \equiv 3 \pmod{4}, \quad 137 \equiv 1 \pmod{4}$$

By quadratic reciprocity,  $\left(\frac{7}{137}\right) = \left(\frac{137}{7}\right)$

$$137 = 7 \cdot 19 + 4 \Rightarrow \left(\frac{137}{7}\right) = \left(\frac{4}{7}\right)$$

Notice that  $4 = 2^2 \equiv 2^2 \pmod{7} \Rightarrow \left(\frac{4}{7}\right) = 1$

This implies:  $\left(\frac{7}{137}\right) = 1$ .

Therefore:

$$\begin{aligned}\left(\frac{-14}{137}\right) &= \left(\frac{-1}{137}\right) \left(\frac{2}{137}\right) \left(\frac{7}{137}\right) \\ &= 1 \cdot 1 \cdot 1 = 1\end{aligned}$$

□

Example: Find  $\left(\frac{55}{179}\right)$

$$\left(\frac{55}{179}\right) = \left(\frac{5}{179}\right) \cdot \left(\frac{11}{179}\right)$$

$$\begin{aligned}5 &\equiv 1 \pmod{4} \\ 11 &\equiv 3 \pmod{4} \\ 179 &\equiv 3 \pmod{4}\end{aligned}$$

$$= \left(\frac{179}{5}\right) \cdot (-1) \cdot \left(\frac{179}{11}\right)$$

$$179 = 5 \cdot 35 + 4 \quad \left(\frac{179}{5}\right) = \left(\frac{4}{5}\right) = 1$$

$$179 = 11 \cdot 16 + 3 \quad \left(\frac{179}{11}\right) = \left(\frac{3}{11}\right) \quad \begin{aligned}3 &\equiv 3 \pmod{4} \\ 11 &\equiv 3 \pmod{4}\end{aligned}$$

$$\left(\frac{3}{11}\right) = -\left(\frac{11}{3}\right) = -\left(\frac{2}{3}\right) = 1 \quad 11 = 3 \cdot 3 + 2$$

$$\Rightarrow \left(\frac{55}{179}\right) = \left(\frac{179}{5}\right) \cdot (-1) \cdot \left(\frac{179}{11}\right)$$

$$= 1 \cdot (-1) \cdot 1 = -1$$

□

Moreover, let  $a, b$  be odd positive integers.

We can write  $a = p_1 p_2 \dots p_r$

$$b = q_1 q_2 \dots q_s$$

Indeed,  $a$  can  
take value  
-1 or 2

We define the Jacobi symbol:

$$\left(\frac{a}{b}\right) = \left(\frac{a}{q_1}\right) \left(\frac{a}{q_2}\right) \dots \left(\frac{a}{q_s}\right)$$

$$= \left(\frac{p_1}{q_1}\right) \left(\frac{p_2}{q_1}\right) \dots \left(\frac{p_r}{q_1}\right) \dots \left(\frac{p_r}{q_s}\right)$$

$$= \prod_{\substack{1 \leq i \leq r \\ 1 \leq j \leq s}} \left(\frac{p_i}{q_j}\right)$$

# Theorem 22.2 (Generalized Law of Quadratic reciprocity)

Let  $a, b$  be odd positive integers,

$$\left(\frac{-1}{b}\right) = \begin{cases} 1 & \text{if } b \equiv 1 \pmod{4} \\ -1 & \text{if } b \equiv 3 \pmod{4} \end{cases}$$

$$\left(\frac{2}{b}\right) = \begin{cases} 1 & \text{if } b \equiv 1, 7 \pmod{8} \\ -1 & \text{if } b \equiv 3, 5 \pmod{8} \end{cases}$$

$$\left(\frac{a}{b}\right) \cdot \left(\frac{b}{a}\right) = (-1)^{\frac{a-1}{2} \cdot \frac{b-1}{2}}$$

Remark: the proof is an application of the original version.

Here we only do one simple example:

$$\left(\frac{-1}{b}\right) = \begin{cases} 1 & \text{if } b \equiv 1 \pmod{4} \\ -1 & \text{if } b \equiv 3 \pmod{4} \end{cases}$$

Proof: We write  $b$  as its primary decomposition

Furthermore, we rearrange the primes of  $b$  such that:

$$b = \underbrace{p_1 p_2 \dots p_r}_{p_i \equiv 1 \pmod{4}} \underbrace{q_1 \dots q_s}_{q_i \equiv 3 \pmod{4}}$$

Exercise: if  $s$  is even, then  $b \equiv 1 \pmod{4}$   
if  $s$  is odd, then  $b \equiv 3 \pmod{4}$ .

Case I:  $s$  is even  $\Rightarrow b \equiv 1 \pmod{4}$

$$\begin{aligned}\left(\frac{-1}{b}\right) &= \left(\frac{-1}{p_1}\right) \cdots \left(\frac{-1}{p_r}\right) \cdot \left(\frac{-1}{q_1}\right) \cdots \left(\frac{-1}{q_s}\right) \\ &= (1)^r \cdot (-1)^s = 1 \quad (s \text{ even}) \\ &= (-1)^{\frac{b-1}{2}} \quad (b \equiv 1 \pmod{4})\end{aligned}$$

Case II:  $s$  is odd  $\Rightarrow b \equiv 3 \pmod{4}$

$$\begin{aligned}\left(\frac{-1}{b}\right) &= \left(\frac{-1}{p_1}\right) \cdots \left(\frac{-1}{p_r}\right) \cdot \left(\frac{-1}{q_1}\right) \cdots \left(\frac{-1}{q_s}\right) \\ &= (1)^r \cdot (-1)^s = -1 \quad (s \text{ odd}) \\ &= (-1)^{\frac{b-1}{2}} \quad (b \equiv 3 \pmod{4})\end{aligned}$$

□.

History: (1) Euler and Lagrange were the first to formulate the Law of Quadratic Reciprocity.

(2) Gauss gave the first proof in 1801.

During his life, he found 7 different proofs.

(3) We will give a proof due to Eisenstein.

(4) The quadratic reciprocity was subsumed into

the Class Field Theory developed by Hilbert, Artin and others from 1890s through 1920s and 1930s.

(5) During 1960s and 1970s, a number of mathematicians formulated a series of conjectures that vastly generalize Class Field Theory and that today go by the name of the Langlands Program.

- The fundamental theorem proved by Wiles in 1995 is a small piece of the Langlands Program, yet it sufficed to solve Fermat's 350-year-old "Last Theorem."